



## Assessment of HMGA2 and PLAG1 rearrangements in breast adenomyoepitheliomas

Pareja, Fresia ; Geyer, Felipe C ; Brown, David N ; Sebastião, Ana P Martins ; Gularte-Mérida, Rodrigo ; Li, Anqi ; Edelweiss, Marcia ; Da Cruz Paula, Arnaud ; Selenica, Pier ; Wen, Hannah Y ; Jungbluth, Achim A ; Varga, Zsuzsanna ; Palazzo, Juan ; Rubin, Brian P ; Ellis, Ian O ; Brogi, Edi ; Rakha, Emad A ; Weigelt, Britta ; Reis-Filho, Jorge S

**Abstract:** Breast adenomyoepitheliomas (AMEs) are rare epithelial-myoepithelial neoplasms that may occasionally produce myxochondroid matrix, akin to pleomorphic adenomas (PAs). Regardless of their anatomic location, PAs often harbor rearrangements involving or . We have recently shown that the repertoire of somatic genetic alterations of AMEs varies according to their estrogen receptor (ER) status; whilst the majority of ER-positive AMEs display mutually exclusive or hotspot mutations, up to 60% of ER-negative AMEs harbor concurrent Q61 hotspot mutations and mutations affecting either or . Here, we hypothesized that a subset of AMEs lacking these somatic genetic alterations could be underpinned by oncogenic fusion genes, in particular those involving or . Therefore, we subjected 13 AMEs to RNA-sequencing for fusion discovery ( = 5) and/or fluorescence in situ hybridization (FISH) analysis for and rearrangements ( = 13). RNA-sequencing revealed an - fusion gene in an ER-positive AME lacking and somatic mutations. This fusion gene, which has been previously described in salivary gland PAs, results in a chimeric transcript composed of exons 1-5 of and exons 3-10 of . No additional in-frame fusion genes or rearrangements were identified in the remaining AMEs analyzed. Our results demonstrate that a subset of AMEs lacking mutations affecting and PI3K pathway-related genes may harbor - fusion genes, suggesting that a subset of breast AMEs may be genetically related to PAs or that a subset of AMEs may originate in the context of a PA.

DOI: <https://doi.org/10.1038/s41523-018-0101-7>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-165678>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Pareja, Fresia ; Geyer, Felipe C ; Brown, David N ; Sebastião, Ana P Martins ; Gularte-Mérida, Rodrigo ; Li, Anqi ; Edelweiss, Marcia ; Da Cruz Paula, Arnaud ; Selenica, Pier ; Wen, Hannah Y ; Jungbluth, Achim A ; Varga, Zsuzsanna ; Palazzo, Juan ; Rubin, Brian P ; Ellis, Ian O ; Brogi, Edi ; Rakha, Emad A ; Weigelt,

Britta; Reis-Filho, Jorge S (2019). Assessment of HMGA2 and PLAG1 rearrangements in breast adenomyoepitheliomas. *npj Breast Cancer*, 5:6.  
DOI: <https://doi.org/10.1038/s41523-018-0101-7>

## ARTICLE OPEN

Assessment of *HMGA2* and *PLAG1* rearrangements in breast adenomyoepitheliomas

Fresia Pareja<sup>1</sup>, Felipe C. Geyer<sup>1</sup>, David N. Brown<sup>1</sup>, Ana P. Martins Sebastião<sup>1</sup>, Rodrigo Gularte-Mérida<sup>1</sup>, Anqi Li<sup>1</sup>, Marcia Edelweiss<sup>1</sup>, Arnaud Da Cruz Paula<sup>1</sup>, Pier Selenica<sup>1</sup>, Hannah Y. Wen<sup>1</sup>, Achim A. Jungbluth<sup>1</sup>, Zsuzsanna Varga<sup>2</sup>, Juan Palazzo<sup>3</sup>, Brian P. Rubin<sup>4</sup>, Ian O. Ellis<sup>5</sup>, Edi Brogi<sup>1</sup>, Emad A. Rakha<sup>5</sup>, Britta Weigelt<sup>1</sup> and Jorge S. Reis-Filho<sup>1</sup>

Breast adenomyoepitheliomas (AMEs) are rare epithelial-myoepithelial neoplasms that may occasionally produce myxochondroid matrix, akin to pleomorphic adenomas (PAs). Regardless of their anatomic location, PAs often harbor rearrangements involving *HMGA2* or *PLAG1*. We have recently shown that the repertoire of somatic genetic alterations of AMEs varies according to their estrogen receptor (ER) status; whilst the majority of ER-positive AMEs display mutually exclusive *PIK3CA* or *AKT1* hotspot mutations, up to 60% of ER-negative AMEs harbor concurrent *HRAS* Q61 hotspot mutations and mutations affecting either *PIK3CA* or *PIK3R1*. Here, we hypothesized that a subset of AMEs lacking these somatic genetic alterations could be underpinned by oncogenic fusion genes, in particular those involving *HMGA2* or *PLAG1*. Therefore, we subjected 13 AMEs to RNA-sequencing for fusion discovery ( $n = 5$ ) and/or fluorescence in situ hybridization (FISH) analysis for *HMGA2* and *PLAG1* rearrangements ( $n = 13$ ). RNA-sequencing revealed an *HMGA2-WIF1* fusion gene in an ER-positive AME lacking *HRAS*, *PIK3CA* and *AKT1* somatic mutations. This fusion gene, which has been previously described in salivary gland PAs, results in a chimeric transcript composed of exons 1–5 of *HMGA2* and exons 3–10 of *WIF1*. No additional in-frame fusion genes or *HMGA2* or *PLAG1* rearrangements were identified in the remaining AMEs analyzed. Our results demonstrate that a subset of AMEs lacking mutations affecting *HRAS* and PI3K pathway-related genes may harbor *HMGA2-WIF1* fusion genes, suggesting that a subset of breast AMEs may be genetically related to PAs or that a subset of AMEs may originate in the context of a PA.

npj Breast Cancer (2019)5:6; <https://doi.org/10.1038/s41523-018-0101-7>

## INTRODUCTION

Breast adenomyoepitheliomas (AMEs) are rare neoplasms with dual epithelial-myoepithelial differentiation,<sup>1</sup> composed of gland-like structures containing an inner layer of pink, eosinophilic epithelial cells and an abluminal layer of often clear, myoepithelial cells. AMEs can display a variety of histologic appearances, and be either estrogen receptor (ER)-positive or ER-negative.<sup>1,2</sup> Although there is overlap in the histologic features of ER-positive and ER-negative AMEs, we have recently shown that the repertoire of genetic alterations of these tumors vary according to their ER status.<sup>3</sup> Whilst ER-negative AMEs harbor *HRAS* Q61 hotspot mutations co-occurring with mutations affecting *PIK3CA* or *PIK3R1* in up to 60% of cases, the majority of ER-positive AMEs were found to display seemingly mutually exclusive *PIK3CA* or *AKT1* activating hotspot mutations.<sup>3</sup>

In the spectrum of histologic appearances of AMEs, myxochondroid matrix has been occasionally described.<sup>4</sup> This type of matrix bears histologic resemblance to the matrix of pleomorphic adenomas (PAs),<sup>1</sup> epithelial-myoepithelial neoplasms that may arise in various anatomic locations, including the breast.<sup>5</sup> PAs are underpinned by recurrent gene rearrangements involving *HMGA2* or *PLAG1* in up to 65% of cases, regardless of their anatomic origin.<sup>6–8</sup> Due to the overlapping histologic appearances of AMEs

and PAs, we sought to determine whether a subset of AMEs, primarily those lacking mutations affecting known drivers (e.g., *HRAS* or PI3K pathway-related genes), would be genetically related to PAs, and would be underpinned by fusion genes, in particular those involving *HMGA2* and *PLAG1*.

## RESULTS

*HMGA2-WIF1* fusion gene in an ER-positive AME

Thirteen breast AMEs, whose whole-exome, targeted capture and/or Sanger sequencing and ER status were previously described in Geyer et al.,<sup>3</sup> were included in this study (Table 1). Six cases were ER-negative and seven were ER-positive. Four ER-negative AMEs harbored concurrent *PIK3CA* and *HRAS* mutations (4/6), one harbored an *HRAS* Q61K mutation and concurrent likely pathogenic *PIK3R1* mutations (1/6), and one was *HRAS* wild-type and harbored a *PIK3CA* mutation (1/6). Five ER-positive AMEs harbored *PIK3CA* mutations (5/7), and all were wild-type for *HRAS*. None of the cases harbored mutations affecting the *AKT1* E17 hotspot locus (Fig. 1a and Supplementary Fig. 1). Notably, all *HRAS* and *PIK3CA* mutations were classical activating hotspot mutations, except for one *PIK3CA* mutation (Q546) which targeted a hotspot residue and was predicted to be likely pathogenic (Fig. 1a).

<sup>1</sup>Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY, USA; <sup>2</sup>Institute of Surgical Pathology, University Hospital Zurich, Zurich, Switzerland;

<sup>3</sup>Department of Pathology, Thomas Jefferson University Hospital, Philadelphia, PA, USA; <sup>4</sup>Department of Pathology, Cleveland Clinic, Cleveland, OH, USA and <sup>5</sup>Department of Pathology, University of Nottingham, Nottingham, UK

Correspondence: Fresia Pareja ([parejaf@mskcc.org](mailto:parejaf@mskcc.org)) or Jorge S. Reis-Filho ([reisfilj@mskcc.org](mailto:reisfilj@mskcc.org))

These authors contributed equally: Fresia Pareja, Felipe C. Geyer

Received: 12 September 2018 Accepted: 27 November 2018

Published online: 18 January 2019

**Table 1.** Clinicopathological and selected genetic features of the 13 breast adenomyoepitheliomas included in this study

Case ID	ER status	PIK3CA status	AKT1 status	PIK3R1 status	HRAS status	Oncogenic fusion genes by RNA sequencing	PLAG1/ HMGA2 rearrangement by FISH
AM1	Negative	MUT	WT	WT	MUT	NT	Negative
AM4	Negative	MUT	WT	WT	MUT	NT	Negative
AM8	Negative	MUT	WT	WT	MUT	NT	Negative
AM11	Negative	MUT	WT	NT	MUT	NT	Negative
AM5	Negative	WT	WT	MUT	MUT	NT	Negative
AM7	Negative	MUT	WT	WT	WT	None	Negative
AM2	Positive	MUT	WT	WT	WT	None	Negative
AM3	Positive	MUT	WT	WT	WT	None	Negative
AM12	Positive	MUT	WT	NT	WT	NT	Negative
AM13	Positive	MUT	WT	NT	WT	NT	Negative
AM17	Positive	MUT	WT	NT	WT	NT	Negative
AM16	Positive	WT	WT	NT	WT	HMGA2-WIF1	Positive (HMGA2)
AM6	Positive	WT	WT	WT	WT	None	Negative

ER estrogen receptor, MUT mutant, NT not tested, WT wild-type

To determine whether AMEs lacking *HRAS* Q61 hotspot mutations would harbor fusion genes, we subjected five *HRAS* wild-type AMEs with available material to RNA-sequencing analysis for an unbiased detection of expressed fusion genes (Supplementary Fig. 1). Using a validated pipeline for the *de novo* discovery of fusion genes,<sup>9</sup> we identified an *HMGA2-WIF1* fusion gene in an ER-positive *HRAS*-/PIK3CA-/AKT1-wild-type AME (AM16) (Figs. 1a–c, Table 1 and Supplementary Table 1). The *HMGA2-WIF1* fusion gene identified in AM16 results in a chimeric transcript encompassing all five exons and the initial segment of the 3' UTR of *HMGA2* fused to exons 3–10 of *WIF1*, and is predicted to be translated to a full length HMGA2 protein and an N-terminal truncated WIF1 protein, with a truncated WIF domain. *WIF1* encodes for a tumor suppressor that modulates Wnt signaling, a role that requires an intact WIF domain.<sup>10</sup>

The AME found to harbor the *HMGA2-WIF1* fusion gene (AM16) displayed the typical histologic features of AMEs,<sup>3</sup> and constituted a well-circumscribed lesion with pushing borders, surrounded by a thick fibrous capsule. This lesion displayed a nodular architecture and a mixed tubular and papillary growth pattern. No cellular atypia, mitotic activity or necrosis was identified (Fig. 1c). Focal areas with conspicuous stroma with myxoid quality were observed (Fig. 1c). The myoepithelial component was highlighted by p63 on immunohistochemical analysis, and strong ER expression was observed in the epithelial component (Fig. 1c).

Given that in the salivary glands, epithelial-myoepithelial carcinomas, the salivary gland counterpart of breast AME, can occasionally originate in the context of a PA (i.e., the so-called carcinoma ex-PA),<sup>11</sup> we sought to define if AM16 would have areas diagnostic of PA. An independent pathology review of all slides available from this AME by five pathologists failed to reveal any areas that would be consistent with a diagnosis of PA.

#### Breast AMEs lack recurrent *HMGA2* or *PLAG1* rearrangements

None of the additional AMEs subjected to RNA-sequencing harbored other fusion genes involving gene partners previously described in PAs,<sup>12</sup> in myoepitheliomas of other anatomical sites (i.e., *EWSR1* and *FUS* rearrangements),<sup>13,14</sup> or in other tumors displaying myoepithelial differentiation (i.e., *CRTC1-MAML2* fusion gene in mucoepidermoid carcinomas or *MYB* and *MYBL1* rearrangements in adenoid cystic carcinoma).<sup>9,15</sup> No additional

likely pathogenic in-frame fusion gene was identified in the cases subjected to RNA-sequencing analysis (Supplementary Table 1).


Given that *HMGA2* and *PLAG1* rearrangements have been described in other neoplasms with epithelial-myoepithelial differentiation, in particular in PAs, we sought to define whether AMEs may harbor fusion genes known to underpin PAs. We subjected the five AMEs analyzed by RNA-sequencing and all the other AMEs included in this study (n = 8) to FISH using *HMGA2* and *PLAG1* dual-color break apart probes (Table 1 and Supplementary Fig. 1). This analysis confirmed the presence of an *HMGA2* rearrangement in both the epithelial and myoepithelial components of AM16 (Fig. 2b) and did not reveal any additional AMEs harboring *HMGA2* or *PLAG1* rearrangements (Table 1).

#### DISCUSSION

We have previously shown that approximately 60% ER-positive AMEs harbor *PIK3CA* or *AKT1* mutations, whereas up to 60% of ER-negative AMEs are characterized by *HRAS* Q61 mutations concurrent with mutations affecting genes of the PI3K signaling pathway.<sup>3</sup> Given the occasional histologic similarities between AMEs and PAs, and the fact that a subset of AMEs lack a known driver genetic alteration in the form of somatic mutations affecting protein coding genes, we posited that a subset of AMEs may harbor oncogenic fusion genes previously described in other myoepithelial lesions including PAs. Our analyses resulted in the identification of an ER-positive *HRAS*-/PIK3CA-/AKT1-wild type AME harboring an *HMGA2-WIF1* fusion gene, which has been described in PAs and carcinomas ex-PA of the salivary gland.<sup>10,16</sup>

The high-mobility group AT-hook 2 (*HMGA2*) gene encodes for a transcriptional regulator of genes involved in cell proliferation and cell death.<sup>17</sup> *HMGA2* overexpression plays a key role in oncogenic transformation through several mechanisms, such as the induction of E2F1 and AP1 activity, promotion of cyclin A expression, inactivation of p53-dependent apoptosis, and activation of the TGF- $\beta$  signaling pathway.<sup>17,18</sup> The Wnt signaling pathway is regulated by secreted antagonists that bind to Wnt proteins, preventing ligand-receptor interactions,<sup>19</sup> such as WIF1.<sup>20</sup> WIF1 consists of an N-terminal secretion signal, five EGF-like domains, a hydrophilic C-terminus, and a WIF domain, which is required for binding to Wnt proteins and for the tumor suppressor properties of WIF1.<sup>19</sup> The *HMGA2-WIF1* fusion gene identified in AM16 has been shown to result in increased HMGA2 expression, presumably






**Fig. 1** Fusion genes involving *HMGA2* or *PLAG1* and somatic mutations targeting *HRAS*, *PIK3CA*, *AKT1* and *PIK3R1* in breast adenomyoepitheliomas. **a** Heatmap depicting fusion gene and somatic mutations targeting *HRAS* Q61, *PIK3CA* and *AKT* E17 hotspot loci and *PIK3R1* mutations identified in breast adenomyoepitheliomas (AMEs;  $n = 13$ ). Cases are shown in columns and genes in rows. Hotspot mutations are annotated as per Chang et al.<sup>23</sup> **b** Representative Sanger sequencing electropherograms of *HRAS* Q61 and *PIK3CA* hotspot loci in AM16. **c** Representative hematoxylin and eosin micrographs of an AME harboring an *HMGA2-WIF1* fusion gene (AM16), and micrographs depicting p63 and estrogen receptor expression. Scale bars, 500  $\mu$ m (upper left), 100  $\mu$ m (upper right) and 50  $\mu$ m (middle and lower panels). ER estrogen receptor, FISH fluorescence in situ hybridization, SNV single nucleotide variant, WT wild-type

due to loss of regulatory sites in its 3' UTR,<sup>21</sup> and decreased *WIF1* expression.<sup>10</sup> The *HMGA2-WIF1* chimeric transcript identified in our study is predicted to encode a full length *HMGA2* protein and an N-terminally truncated *WIF1* protein harboring a truncated *WIF* domain. Given that the *HMGA2* breakpoint maps to its 3' UTR after the stop codon, it is possible that *WIF1* may not even be translated, akin to a rearrangement involving the same *HMGA2* and *WIF1* exons previously described in a PA arising in the salivary gland.<sup>16</sup> In addition, *HMGA2* and *WIF1* display opposite transcriptional orientations, and this fusion gene may stem from a cryptic paracentric inversion (Fig. 2a).<sup>16</sup>

Taken together, the *HMGA2-WIF1* fusion identified here might result in increased expression of *HMGA2*, with ensuing activation of TGF- $\beta$  signaling, along with derepression of Wnt signaling.

Taken together, our findings suggest that a subset of AMEs lacking genetic alterations involving genes of the RAS-MAPK pathway may be underpinned by fusion genes resulting in the activation of alternative signaling pathways, such as TGF- $\beta$  and Wnt.

One could posit that the AME harboring the *HMGA2-WIF1* fusion gene described in this study would, in fact, constitute a breast PA. This case was independently reviewed by five breast pathologists who concurred in the diagnosis of AME. It should be noted, however, that despite being a *bona fide* AME, this case focally displayed increased myxoid stroma, bearing some resemblance to, but not fulfilling the diagnostic criteria for, a breast PA. Another potential explanation for the presence of this fusion gene in AM16 is that it would constitute the breast equivalent of the salivary gland epithelial-myoepithelial carcinoma ex-PA.<sup>11</sup> The central



Our study was several limitations, such as the small size of our cohort and the fact that we included only two *HRAS*-/*PIK3CA*-/*AKT1*-wild type AMEs, given that no additional material from other AMEs was available for transcriptomic or FISH analysis. The limited sample size of our study precludes definitive conclusions regarding the relationship between ER status and the presence of the *HMGA2-WIF1* fusion gene in AMEs to be drawn. Despite these limitations, our findings demonstrate that AMEs lacking mutations affecting *HRAS*, *PIK3CA* and *AKT1* may harbor the *HMGA2-WIF1* fusion gene, previously described in salivary gland PAs and carcinomas ex-PA,<sup>10,16</sup> suggesting that a subset of AMEs may be genetically related to PAs or that AMEs may originate in the context of breast PAs.

RNA-sequencing was performed on five *HRAS*-wild type AMEs according to standard protocols employed at the Integrated Genomics Operation of Memorial Sloan Kettering Cancer Center (MSKCC).<sup>27</sup> In brief, paired-end massively parallel RNA-sequencing (2 × 50 bp) was performed on a HiSeq2000 (Illumina), as previously described.<sup>28</sup> Read pairs supporting chimeric transcripts were identified using deFuse,<sup>29</sup> and INTEGRATE,<sup>30</sup> followed by exclusion of candidate fusion transcripts found in a set of 287 normal breast tissues from the TCGA dataset,<sup>31</sup> as previously described.<sup>28</sup> The Bayesian probability of the remaining candidate fusion genes, supported by at least two spanning reads, to constitute drivers was annotated using OncoFuse<sup>32</sup> as previously described.<sup>28</sup>



## Fluorescence in situ hybridization (FISH)

All cases included in this study ( $n = 13$ ) were subjected to FISH analysis for *HMGA2* and *PLGA1* using dual-color break-apart probes following validated protocols at the MSKCC Molecular Cytogenetics Core, as previously described.<sup>33</sup> The probe mix consisted of bacterial artificial chromosome (BAC) clones mapping to 5' *HMGA2* (RP11-230G5, RP11-662G15; red) and 3' *HMGA2* (RP11-937C6, RP11-167E10; green), and BAC clones mapping to 5' *PLAG1* (RP11-92A9, RP11-111I18; red) and 3' *PLAG1* (RP11-144E19, RP11-246A9; green). A minimum of 50 interphase nuclei were analyzed for *HMGA2* or *PLAG1* rearrangements. Cases were considered positive for rearrangement if separation of the 5' (red) and 3' (green) signals ( $>2$  signal width apart) was identified in  $>15\%$  tumor cells. FISH analyses were performed with observers blinded to the results of the RNA-sequencing analysis.

## DATA AVAILABILITY

RNA-sequencing data that support the findings of this study have been deposited in the NCBI Sequence Read Archive (SRA) under the accession code SRP158271. WES and MSK-IMPACT sequencing data retrieved from Geyer et al.<sup>3</sup> are available in SRA under the accession numbers SRP065277 and SRP065302, respectively. All available data are available from the authors.

## ACKNOWLEDGEMENTS

This study was funded by the Breast Cancer Research Foundation. Research reported in this publication was funded in part by a Cancer Center Support Grant of the National Institutes of Health/National Cancer Institute (grant No P30CA008748). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## AUTHORS' CONTRIBUTIONS

B.W. and J.S.R.-F. conceived the study. H.Y.W., Z.V., J.P., B.P.R., I.O.E., E.B. and E.A.R. provided tissue samples. F.P., F.C.G., M.E., A.P.M.S., E.A.R., I.O.E. and J.S.R.-F. performed the pathology review. Data acquisition, interpretation and analysis were performed by F.P., F.C.G., D.N.B., A.L., A.P.M.S., R.G.-M., A.d.C.P., P.S. and A.A.J. The manuscript was initially drafted by F.P. and J.S.R.-F., and all authors edited and approved the final draft of the manuscript. F.P. and F.C.G. contributed equally to this work.

## ADDITIONAL INFORMATION

**Supplementary information** accompanies the paper on the *npj Breast Cancer* website (<https://doi.org/10.1038/s41523-018-0101-7>).

**Competing interests:** J.S.R.-F. reports personal/consultancy fees from VolitionRx, Page.AI, Goldman Sachs, Grail, Ventana Medical Systems and Genentech, outside the submitted work. The remaining authors declare no competing interests.

**Publisher's note:** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## REFERENCES

- Yoon, J. Y. & Chitale, D. Adenomyoepithelioma of the breast: a brief diagnostic review. *Arch. Pathol. Lab. Med.* **137**, 725–729 (2013).
- Lakhani, S. R., Ellis, I. O., Schnitt, S. J., Tan, P. H., van de Vijver, M. J. *WHO Classification of Breast Tumors*. (International Agency for Research Cancer (IARC), Lyon, 2012).
- Geyer, F. C. et al. Recurrent hotspot mutations in HRAS Q61 and PI3K-AKT pathway genes as drivers of breast adenomyoepitheliomas. *Nat. Commun.* **9**, 1816 (2018).
- McLaren, B. K., Smith, J., Schuyler, P. A., Dupont, W. D. & Page, D. L. Adenomyoepithelioma: clinical, histologic, and immunohistologic evaluation of a series of related lesions. *Am. J. Surg. Pathol.* **29**, 1294–1299 (2005).
- Foschini, M. P. et al. The morphological spectrum of salivary gland type tumours of the breast. *Pathology* **49**, 215–227 (2017).
- Stenman, G. Fusion oncogenes in salivary gland tumors: molecular and clinical consequences. *Head. Neck Pathol.* **7**(Suppl 1), S12–S19 (2013).
- Bahrami, A., Dalton, J. D., Krane, J. F. & Fletcher, C. D. A subset of cutaneous and soft tissue mixed tumors are genetically linked to their salivary gland counterpart. *Genes Chromosomes Cancer* **51**, 140–148 (2012).
- Pareja, F. et al. Pleomorphic adenoma and mucoepidermoid carcinoma of the breast are underpinned by fusion genes. *Mod. Pathol.* **31**, 98–98 (2018).

- Kim, J. et al. MYBL1 rearrangements and MYB amplification in breast adenoid cystic carcinomas lacking the MYB-NFIB fusion gene. *J. Pathol.* **244**, 143–150 (2018).
- Queimado, L., Lopes, C. S. & Reis, A. M. WIF1, an inhibitor of the Wnt pathway, is rearranged in salivary gland tumors. *Genes Chromosomes Cancer* **46**, 215–225 (2007).
- Antony, J., Gopalan, V., Smith, R. A. & Lam, A. K. Carcinoma ex pleomorphic adenoma: a comprehensive review of clinical, pathological and molecular data. *Head. Neck Pathol.* **6**, 1–9 (2012).
- Antonescu, C. R. et al. Frequent PLAG1 gene rearrangements in skin and soft tissue myoepithelioma with ductal differentiation. *Genes Chromosomes Cancer* **52**, 675–682 (2013).
- Huang, S. C. et al. Novel FUS-KLF17 and EWSR1-KLF17 fusions in myoepithelial tumors. *Genes Chromosomes Cancer* **54**, 267–275 (2015).
- Antonescu, C. R. et al. EWSR1-POU5F1 fusion in soft tissue myoepithelial tumors. A molecular analysis of sixty-six cases, including soft tissue, bone, and visceral lesions, showing common involvement of the EWSR1 gene. *Genes Chromosomes Cancer* **49**, 1114–1124 (2010).
- Tonon, G. et al. t(11;19)(q21; p13) translocation in mucoepidermoid carcinoma creates a novel fusion product that disrupts a Notch signaling pathway. *Nat. Genet.* **33**, 208–213 (2003).
- Persson, F. et al. High-resolution genomic profiling of adenomas and carcinomas of the salivary glands reveals amplification, rearrangement, and fusion of HMGA2. *Genes Chromosomes Cancer* **48**, 69–82 (2009).
- Fusco, A. & Fedele, M. Roles of HMGA proteins in cancer. *Nat. Rev. Cancer* **7**, 899–910 (2007).
- Morishita, A. et al. HMGA2 is a driver of tumor metastasis. *Cancer Res* **73**, 4289–4299 (2013).
- Malinauskas, T., Aricescu, A. R., Lu, W., Siebold, C. & Jones, E. Y. Modular mechanism of Wnt signaling inhibition by Wnt inhibitory factor 1. *Nat. Struct. Mol. Biol.* **18**, 886–893 (2011).
- Hsieh, J. C. et al. A new secreted protein that binds to Wnt proteins and inhibits their activities. *Nature* **398**, 431–436 (1999).
- Mayr, C., Hemann, M. T. & Bartel, D. P. Disrupting the pairing between let-7 and Hmga2 enhances oncogenic transformation. *Science* **315**, 1576–1579 (2007).
- Hammond, M. E. et al. American Society of Clinical Oncology/College Of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. *J. Clin. Oncol.* **28**, 2784–2795 (2010).
- Chang, M. T. et al. Identifying recurrent mutations in cancer reveals widespread lineage diversity and mutational specificity. *Nat. Biotechnol.* **34**, 155–163 (2016).
- Martelotto, L. G. et al. Genomic landscape of adenoid cystic carcinoma of the breast. *J. Pathol.* **237**, 179–189 (2015).
- Wetterskog, D. et al. Adenoid cystic carcinomas constitute a genomically distinct subgroup of triple-negative and basal-like breast cancers. *J. Pathol.* **226**, 84–96 (2012).
- Ashworth, A., Lord, C. J. & Reis-Filho, J. S. Genetic interactions in cancer progression and treatment. *Cell* **145**, 30–38 (2011).
- Weinreb, I. et al. Hotspot activating PRKD1 somatic mutations in polymorphous low-grade adenocarcinomas of the salivary glands. *Nat. Genet.* **46**, 1166–1169 (2014).
- Piscuoglio, S. et al. Integrative genomic and transcriptomic characterization of papillary carcinomas of the breast. *Mol. Oncol.* **8**, 1588–1602 (2014).
- McPherson, A. et al. deFuse: an algorithm for gene fusion discovery in tumor RNA-Seq data. *PLoS Comput. Biol.* **7**, e1001138 (2011).
- Zhang, J. et al. INTEGRATE: gene fusion discovery using whole genome and transcriptome data. *Genome Res.* **26**, 108–118 (2015).
- Cancer Genome Atlas N. Comprehensive molecular portraits of human breast tumours. *Nature* **490**, 61–70 (2012).
- Shugay, M., Ortiz de Mendibil, I., Vizmanos, J. L. & Novo, F. J. Oncofuse: a computational framework for the prediction of the oncogenic potential of gene fusions. *Bioinformatics* **29**, 2539–2546 (2013).
- Piscuoglio, S. et al. Uterine adenocarcinomas are mesenchymal neoplasms. *J. Pathol.* **238**, 381–388 (2016).



**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019